

# HIGHER EVEN DIMENSIONAL REIDEMEISTER TORSION FOR TORUS KNOT EXTERIORS

YOSHIKAZU YAMAGUCHI

**ABSTRACT.** We study that the asymptotic of higher dimensional Reidemeister torsion for torus knot exteriors, which are related to the results by W. Müller [Mül] and P. Menal-Ferrer and J. Porti [MFPa] on asymptotics of Reidemeister torsion and hyperbolic volumes for hyperbolic 3-manifolds. We show that the sequence  $\frac{1}{(2N)^2} \log |\text{Tor}(E_K; \rho_{2N})|$  converges to zero when  $N$  goes to infinity where  $\text{Tor}(E_K; \rho_{2N})$  is the higher dimensional Reidemeister torsion of a torus knot exterior and an acyclic  $\text{SL}_{2N}(\mathbb{C})$ -representation of the torus knot group. We also give a complete classification of  $\text{SL}_2(\mathbb{C})$ -representations of torus knot groups, which induce acyclic  $\text{SL}_{2N}(\mathbb{C})$ -representations.

## 1. INTRODUCTION

For an oriented hyperbolic 3-manifold with finite volume, the hyperbolic volume is related to the asymptotic behavior for the sequence of Reidemeister torsion respect to  $\text{SL}_n(\mathbb{C})$ -representations of the fundamental group when  $n$  increases to infinity. Here we are concerned with  $\text{SL}_n(\mathbb{C})$ -representations of fundamental groups defined as the composition of an  $\text{SL}_2(\mathbb{C})$ -representation corresponding to the complete hyperbolic structure and the  $n$ -dimensional irreducible representation of  $\text{SL}_2(\mathbb{C})$ . This result was shown by W. Müller [Mül] in the context of Ray–Singer analytic torsion for closed hyperbolic 3-manifolds and then was extended to the cusped hyperbolic 3-manifolds, according to P. Menal-Ferrer and J. Porti [MFPa].

Menal-Ferrer and Porti take advantage of the combinatorial description of Reidemeister torsion to establish the relation between the asymptotic behavior for a sequence of Reidemeister torsions and the hyperbolic volume of a cusped hyperbolic 3-manifold  $M$ :

$$\lim_{n \rightarrow \infty} \frac{\log |\text{Tor}(M; \rho_n)|}{n^2} = -\frac{\text{Vol}(M)}{4\pi}$$

where  $\text{Tor}(M; \rho_n)$  denotes the Reidemeister torsion respect to  $\text{SL}_n(\mathbb{C})$ -representation  $\rho_n$  of  $\pi_1(M)$ . We will see a little more detail in the below.

We can also take advantage of combinatorial feature of Reidemeister torsion to consider the case for non-hyperbolic manifolds  $M$  and expect that the sequence  $\frac{1}{n^2} \log |\text{Tor}(M; \rho_n)|$  converges to zero when  $n \rightarrow \infty$ . In this paper, we will show that this expected asymptotic hold for all torus knots exteriors and even dimensional *acyclic* representations  $\rho_{2N}$  (see the subsequent of Definition 2.1 for the terminology “acyclic”). To define Reidemeister torsion, we need a homomorphism from fundamental groups of manifolds to some linear group, which are called *representations*. In the case for oriented hyperbolic 3-manifolds  $M$ , we can choose  $\text{SL}_2(\mathbb{C})$ -representations corresponding to the hyperbolic structure, called holonomy representations, since  $M$  is expressed as  $\Gamma \backslash \mathbb{H}^3$  where  $\Gamma$  is a discrete subgroup

*Date:* August 23, 2012.

*2000 Mathematics Subject Classification.* Primary: 57M27, 57M05, 57M12, Secondary: 57M25.

*Key words and phrases.* torus knots; irreducible representations; Reidemeister torsion; hyperbolic volume.

in the orientation preserving isometries  $\text{Isom}^+(\mathbb{H}^3) \simeq \text{SL}_2(\mathbb{C})/\{\pm I\}$ . On the other hand, we have no specific such as holonomy representations for non-hyperbolic 3-manifolds. Hence we need to deal with all representations of torus knot groups to discuss the asymptotic of Reidemeister torsions. However, for the simplicity, we focus on even dimensional acyclic representations of torus knot groups since Raghunathan's cohomology vanishing Theorem [Rag65] and the result of Menal-Ferrer and Porti [MFPb] say that all even dimensional  $\rho_{2N}$  for a lift  $\rho$  of holonomy representation are acyclic. We will establish the following asymptotic behaviors of sequences of Reidemeister torsions for torus knot exteriors respect to acyclic  $\text{SL}_{2N}(\mathbb{C})$ -representations  $\rho_{2N}$ .

**Main Theorem** (Theorem 4.2). *Let  $E_K$  denote  $S^3 \setminus N(K)$  where  $K$  is a torus knot and  $N(K)$  is an open tubular neighbourhood of  $K$ . Suppose that the composition  $\rho_{2N}$  of an irreducible  $\text{SL}_2(\mathbb{C})$ -representation  $\rho$  of  $\pi_1(E_K)$  and the  $2N$ -dimensional irreducible representation of  $\text{SL}_2(\mathbb{C})$  is acyclic. For a torus knot exterior  $E_K$  and  $\rho_{2N}$ , it holds that*

$$\lim_{N \rightarrow \infty} \frac{\log |\text{Tor}(E_K; \rho_{2N})|}{(2N)^2} = 0.$$

Our theorem follows from explicit computations of Reidemeister torsion for torus knot exteriors in Proposition 4.1 and the existence of upper and lower bounds dependently only on  $2N$ . To complete our observation, we have to find out which irreducible  $\text{SL}_2(\mathbb{C})$ -representations of torus knot groups induce acyclic  $\text{SL}_{2N}(\mathbb{C})$ -representations. We will also establish a complete classification which irreducible  $\text{SL}_2(\mathbb{C})$ -representation  $\rho$  gives an acyclic  $\rho_{2N}$  in Proposition 3.1.

## 2. PRELIMINARIES

### 2.1. Reidemeister torsion.

*Torsion for acyclic chain complexes.* *Torsion* is an invariant defined for a based chain complexes. We denote by  $C_*$  the *based* chain complex:

$$C_* : 0 \rightarrow C_n \xrightarrow{\partial_n} C_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \rightarrow 0$$

where each chain module  $C_i$  is a vector space over a field  $\mathbb{F}$  and equipped with a basis  $\mathbf{c}^i$ . We are mainly interested in chain complexes whose homology group vanishes, i.e.,  $H_*(C_*) = \mathbf{0}$ . Such chain complexes are referred to as being *acyclic*. The chain complex  $C_*$  also has a basis determined by the boundary operators  $\partial_i$ , which arises from the following decomposition of chain modules. Roughly speaking, torsion provides a property of a chain complex in the difference between a given basis and new one determined by the boundary operators.

We suppose that a based chain complex  $(C_*, \mathbf{c}^i)$  is acyclic. For each boundary operator  $\partial_i$ , let  $Z_i \subset C_i$  denote the kernel and  $B_i \subset C_{i-1}$  the image by  $\partial_i$ . The chain module  $C_i$  is expressed as the direct sum of  $Z_i$  and the lift of  $B_i$ , denoted by  $\tilde{B}_i$ . Moreover we can rewrite the kernel  $Z_i$  as the image of boundary operator  $\partial_{i+1}$ :

$$\begin{aligned} C_i &= Z_i \oplus \tilde{B}_i \\ &= \partial_{i+1} \tilde{B}_{i+1} \oplus \tilde{B}_i \end{aligned}$$

where  $B_{i+1}$  is written as  $\partial_{i+1} \tilde{B}_{i+1}$ .

We denote by  $\tilde{\mathbf{b}}^i$  a basis of  $\tilde{B}_{i+1}$ . Then the set  $\partial_{i+1}(\tilde{\mathbf{b}}^{i+1}) \cup \tilde{\mathbf{b}}^i$  forms a new basis of the vector space  $C_i$ . We define the *torsion* of  $C_*$  as the following alternating product of

determinants of base change matrices:

$$(1) \quad \text{Tor}(C_*, \mathbf{c}^*) = \prod_{i \geq 0} \left[ \partial_{i+1}(\tilde{\mathbf{b}}^{i+1}) \cup \tilde{\mathbf{b}}^i / \mathbf{c}^i \right]^{(-1)^{i+1}} \in \mathbb{F}^* = \mathbb{F} \setminus \{0\}$$

where  $[\partial_{i+1}(\tilde{\mathbf{b}}^{i+1}) \cup \tilde{\mathbf{b}}^i / \mathbf{c}^i]$  denotes the determinant of base change matrix from the given basis  $\mathbf{c}^i$  to the new one  $\partial_{i+1}(\tilde{\mathbf{b}}^{i+1}) \cup \tilde{\mathbf{b}}^i$ .

Note that the right hand side is independent of the choice of bases  $\tilde{\mathbf{b}}^i$ . The alternating product in (1) is determinant by the based chain complex  $(C_*, \mathbf{c}^*)$ .

*Reidemeister torsion for CW-complexes.* We will consider torsion of *twisted chain complexes* given by a CW-complex and a representation of its fundamental group in this paper. Let  $W$  denote a finite CW-complex and  $(V, \rho)$  a representation of  $\pi_1(W)$ , which means  $V$  is a vector space over  $\mathbb{F}$  and  $\rho$  is a homomorphism from  $\pi_1(W)$  into  $\text{GL}(V)$  which is referred as a  $\text{GL}(V)$ -representation.

**Definition 2.1.** We define the twisted chain complex  $C_*(W; V_\rho)$  which consists of the twisted chain module as:

$$C_i(W; V_\rho) := V \otimes_{\mathbb{Z}[\pi_1(W)]} C_i(\tilde{W}; \mathbb{Z})$$

where  $\tilde{W}$  is the universal cover of  $W$  and  $C_i(\tilde{W}; \mathbb{Z})$  is a left  $\mathbb{Z}[\pi_1(W)]$ -module given by the covering transformation of  $\pi_1(W)$ . In taking the tensor product, we regard  $V$  as a right  $\mathbb{Z}[\pi_1]$ -module under the homomorphism  $\rho^{-1}$ . We identify a chain  $\mathbf{v} \otimes \gamma c$  with  $\rho(\gamma)^{-1}(\mathbf{v}) \otimes c$  in  $C_i(W; V_\rho)$ .

We call  $C_*(W; V_\rho)$  the twisted chain complex with the coefficient  $V_\rho$ . Choosing a basis of the vector space  $V$ , we give a basis of the twisted chain complex  $C_*(W; V_\rho)$ . To be more precise, let  $\{e_1^i, \dots, e_{m_i}^i\}$  be the set of  $i$ -dimensional cells of  $W$  and  $\{\mathbf{v}_1, \dots, \mathbf{v}_d\}$  a basis of  $V$  where  $d = \dim_{\mathbb{F}} V$ . Choosing a lift  $\tilde{e}_j^i$  of each cell and taking tensor product with the basis of  $V$ , we have the following basis of  $C_i(W; V_\rho)$ :

$$\mathbf{c}^i(W; V) = \{\mathbf{v}_1 \otimes \tilde{e}_1^i, \dots, \mathbf{v}_d \otimes \tilde{e}_1^i, \dots, \mathbf{v}_1 \otimes \tilde{e}_{m_i}^i, \dots, \mathbf{v}_d \otimes \tilde{e}_{m_i}^i\}.$$

We denote by  $H_*(W; V_\rho)$  the homology group, which is called *the twisted homology group* and say that  $\rho$  is acyclic if the twisted homology group vanishes. Regarding  $C_*(W; V_\rho)$  as a based chain complex, we define the Reidemeister torsion for  $W$  and an acyclic representation  $(V, \rho)$  as the torsion of  $C_*(W; V_\rho)$ , i.e.,

$$(2) \quad \text{Tor}(W; V_\rho) = \text{Tor}(C_*(W; V_\rho), \mathbf{c}^*(W; V)) \in \mathbb{F}^*$$

up to a factor in  $\{\pm \det(\rho(\gamma)) \mid \gamma \in \pi_1(W)\}$  since we have many choices of lifts  $\tilde{e}_j^i$  and orders and orientations of cells  $e_j^i$ .

Note that the definition (2) does not depend of choice of a basis  $\{\mathbf{v}_1, \dots, \mathbf{v}_d\}$  since the Euler characteristic of  $C_*(W; V_\rho)$  must be zero by the acyclicity. More precisely, the torsion defined by another basis  $\{\mathbf{u}_1, \dots, \mathbf{u}_d\}$  is expressed as the product of the torsion defined by  $\{\mathbf{v}_1, \dots, \mathbf{v}_d\}$  and the following factor:

$$[\{\mathbf{u}_1, \dots, \mathbf{u}_d\} / \{\mathbf{v}_1, \dots, \mathbf{v}_d\}]^{-\chi(W)}.$$

*Remark 2.2.* If we choose an  $\text{SL}(V)$ -representation  $\rho$  with  $\dim_{\mathbb{F}} V$  even, then the torsion  $\text{Tor}(W; V_\rho)$  has no indeterminacy.

See Turaev's book [Tur01] for more detail on Reidemeister torsion.

**2.2. Irreducible representations of  $\mathrm{SL}_2(\mathbb{C})$  and higher dimensional Reidemeister torsion.** We review irreducible representations of  $\mathrm{SL}_2(\mathbb{C})$  briefly. The vector space  $\mathbb{C}^2$  has the standard action of  $\mathrm{SL}_2(\mathbb{C})$  and it is known that the symmetric product  $\mathrm{Sym}^{n-1}(\mathbb{C}^2)$  and the induced action by  $\mathrm{SL}_2(\mathbb{C})$  gives an  $n$ -dimensional irreducible representation of  $\mathrm{SL}_2(\mathbb{C})$ . We can identify  $\mathrm{Sym}^{n-1}(\mathbb{C}^2)$  with the vector space of homogeneous polynomials on  $\mathbb{C}^2$  with degree  $n - 1$ , denoted by  $V_n$ :

$$V_n = \mathrm{span}_{\mathbb{C}} \langle z_1^{n-1}, z_1^{n-2} z_2, \dots, z_1^{n-k-1} z_2^k, \dots, z_1 z_2^{n-2}, z_2^{n-1} \rangle$$

The action of  $A \in \mathrm{SL}_2(\mathbb{C})$  is expressed as

$$(3) \quad A \cdot p(z_1, z_2) = p(A^{-1} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix})$$

where  $p(z_1, z_2)$  is a homogeneous polynomial and the variables in the right hand are determined by the action of  $A^{-1}$  on the column vector by matrix multiplication. We write  $(V_n, \sigma_n)$  for the representation given by the action (3) of  $\mathrm{SL}_2(\mathbb{C})$  where  $\sigma_n$  denotes the homomorphism from  $\mathrm{SL}_2(\mathbb{C})$  into  $\mathrm{GL}(V_n)$ .

*Remark 2.3.* It is known that

- (1) each representation  $(V_n, \sigma_n)$  turns into an irreducible  $\mathrm{SL}_n(\mathbb{C})$ -representation of  $\mathrm{SL}_2(\mathbb{C})$  and;
- (2) every irreducible  $n$ -dimensional representation of  $\mathrm{SL}_2(\mathbb{C})$  is equivalent to  $(V_n, \sigma_n)$ .

It is worth noting that  $\mathrm{SL}_2(\mathbb{C})$  acts on the right in the representation  $(V_n, \sigma_n)$ , relating to our construction of twisted chain complexes in Section 2.1. Let  $\rho$  be a homomorphism from  $\pi_1(W)$  into  $\mathrm{SL}_2(\mathbb{C})$  and  $\rho_n$  the composition  $\sigma_n \circ \rho$ , which gives an  $\mathrm{SL}_n(\mathbb{C})$ -representation of  $\pi_1(W)$ . If the twisted chain complex  $C_*(W; V_n)$  is acyclic, we can consider  $\mathrm{Tor}(W; V_n)$ . We will drop the subscript  $\rho$  in the coefficient for simplicity when no confusion can arise. We define the higher dimensional Reidemeister torsion for  $W$  and  $\rho$  as  $\mathrm{Tor}(W; V_n)$  and denote by  $\mathrm{Tor}(W; \rho_n)$  since the coefficient  $V_n$  of  $C_*(W; V_n)$  is determined by  $\rho$  and  $n$ .

We also review the explicit forms of diagonal matrices under  $\sigma_n$ , which will be needed in our computations. Let  $A \in \mathrm{SL}_2(\mathbb{C})$  be a diagonal matrix  $\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$  and  $p(z_1, z_2)$  the monomial  $z_1^{n-k-1} z_2^k$ . Then the action of  $A$  is expressed as

$$A \cdot p(z_1, z_2) = p(a^{-1} z_1, a z_2) = a^{2k-n+1} z_1^{n-k} z_2^k = a^{2k-n+1} p(z_1, z_2).$$

Hence the eigenvalues of  $\sigma_n(A) \in \mathrm{SL}_n(\mathbb{C})$  are given by  $a^{-n+1}, a^{-n+3}, \dots, a^{n-1}$ , i.e., the weight space of  $\sigma_n$  is  $\{-n + 1, -n + 3, \dots, n - 1\}$  and the multiplicity of each weight is 1.

**2.3.  $\mathrm{SL}_2(\mathbb{C})$ -representations of torus knot groups.** Here and subsequently, let  $K$  denote  $(p, q)$ -torus knot and  $E_K$  the knot exterior. The Reidemeister torsion depends on the conjugacy classes of irreducible representations of fundamental groups. We are interested in irreducible representations from torus knot groups into  $\mathrm{SL}_2(\mathbb{C})$ . We denote by  $R(E_K)$  the set of irreducible  $\mathrm{SL}_2(\mathbb{C})$ -representations of  $\pi_1(E_K)$  and by  $\hat{R}(E_K)$  the set of conjugacy classes of representations in  $R(E_K)$ . According to D. Johnson [Joh], from the presentation  $\langle x, y \mid x^p = y^q \rangle$  of  $\pi_1(E_K)$ , we can regard  $\hat{R}(E_K)$  as a set of punctured complex planes:

**Proposition 2.4** ([Joh], Proposition 3.7 in [KM]). *Let  $K$  be the  $(p, q)$ -torus knot. Then  $\hat{R}(E_K)$  consists of  $(p - 1)(q - 1)/2$  components, which are determined by the following data, denoted by  $\hat{R}_{a,b}(E_K)$ :*

- (1)  $0 < a < p, 0 < b < q$ .
- (2)  $a \equiv b \pmod{2}$ .

- (3) For every  $[\rho] \in \hat{R}_{a,b}(E_K)$ , we have that  $\text{tr } \rho(x) = 2 \cos\left(\frac{\pi a}{p}\right)$  and  $\text{tr } \rho(y) = 2 \cos\left(\frac{\pi b}{q}\right)$ . Moreover irreducible  $\text{SL}_2(\mathbb{C})$ -representation  $\rho$  sends  $x^p (= y^q)$  to  $(-I)^a$ .
- (4)  $\text{tr } \rho(\mu) \neq 2 \cos(\pi(ra/p \pm sb/q))$  where  $\mu$  denotes the meridian given by  $x^{-r}y^s$  and  $r$  and  $s$  are integers satisfying that  $ps - qr = 1$ .

In particular,  $\hat{R}_{a,b}(E_K)$  is parametrized by  $\text{tr } \rho(\mu)$  and the complex dimension one.

Therefore, for every irreducible  $\text{SL}_2(\mathbb{C})$ -representation  $\rho$ , the  $\text{SL}_2(\mathbb{C})$ -matrices  $\rho(x)$  and  $\rho(y)$  are diagonalizable, i.e., they are conjugate to diagonal matrices:

$$\begin{pmatrix} e^{a\pi\sqrt{-1}/p} & 0 \\ 0 & e^{-a\pi\sqrt{-1}/p} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} e^{b\pi\sqrt{-1}/q} & 0 \\ 0 & e^{-b\pi\sqrt{-1}/q} \end{pmatrix}$$

when the conjugacy class of  $\rho$  is contained in  $\hat{R}_{a,b}(E_K)$ . Then the eigenvalues of  $\rho_{2N}(x)$  are  $e^{\pm(2N-1)a\pi\sqrt{-1}/p}$ ,  $e^{\pm(2N-3)a\pi\sqrt{-1}/p}$ ,  $\dots$ ,  $e^{\pm a\pi\sqrt{-1}/p}$  and those of  $\rho_{2N}(y)$  are  $e^{\pm(2N-1)b\pi\sqrt{-1}/q}$ ,  $e^{\pm(2N-3)b\pi\sqrt{-1}/q}$ ,  $\dots$ ,  $e^{\pm b\pi\sqrt{-1}/q}$ .

### 3. TWISTED HOMOLOGY GROUPS OF TORUS KNOT EXTERIORS

We consider the acyclicity for the twisted chain complex of  $E_K$  defined by  $\rho_{2N}$ . The twisted homology group is invariant under homotopy equivalence. We first construct a 2-dimensional CW-complex  $W$  which is homotopy equivalent to a torus knot exterior  $E_K$  and then observe the twisted chain complex  $C_*(W; V_{2N})$  instead of  $C_*(E_K; V_{2N})$ .

From the presentation  $\langle x, y \mid x^p = y^q \rangle$  of  $\pi_1(E_K)$ , we can construct 2-dimensional CW-complex  $W$  consisting of one 0-cell  $e^0$  and two 1-cells  $e_i^1$  ( $i = 1, 2$ ) and one 2-cell  $e^2$  as in Fig. 1. The closed loops  $e^0 \cup e_1^1$  and  $e^0 \cup e_2^1$  correspond to the generators  $x$  and  $y$ .




FIGURE 1. CW-complex  $W$

Let  $\rho$  be an irreducible  $\text{SL}_2(\mathbb{C})$ -representation whose conjugacy class lies in  $\hat{R}_{a,b}(E_K)$ . The composition  $\rho_{2N} = \sigma_{2N} \circ \rho$  also defines the twisted chain complex of  $W$  as follows:

$$(4) \quad 0 \rightarrow C_2(W; V_{2N}) \xrightarrow{\partial_2} C_1(W; V_{2N}) \xrightarrow{\partial_1} C_0(W; V_{2N}) \rightarrow 0$$

where  $V_{2N}$  is the  $2N$ -dimensional vector space over  $\mathbb{C}$ . When we write  $X$  and  $Y$  for the matrices  $\rho_{2N}(x)$  and  $\rho_{2N}(y)$ , the boundary operators  $\partial_i$  ( $i = 1, 2$ ) are expressed as

$$(5) \quad \partial_2 = \begin{pmatrix} I + X + \dots + X^{p-1} \\ -(I + Y + \dots + Y^{q-1}) \end{pmatrix}, \quad \partial_1 = \begin{pmatrix} X - I & Y - I \end{pmatrix}.$$

The acyclicity of  $C_*(W; V_{2N})$  is determined by the parity of parameters  $a$  and  $b$  for the component  $\hat{R}_{a,b}(E_K)$  containing the conjugacy class  $\rho$ , i.e., the acyclicity of  $C_*(E_K; V_{2N})$  depends on only the component  $\hat{R}_{a,b}(E_K)$ .

**Proposition 3.1.** *Suppose that the conjugacy classes of  $\rho$  is contained in  $\hat{R}_{a,b}(E_K)$ .*

- (1) If the integers  $a$  and  $b$  are odd, then  $C_*(E_K; V_{2N})$  is acyclic, i.e.,  $H_*(E_K; V_{2N}) = \mathbf{0}$ .  
(2) If the integers  $a$  and  $b$  are even, then  $C_*(E_K; V_{2N})$  is not acyclic.

*Proof.* It is enough to observe  $C_*(W; V_{2N})$  instead of  $C_*(E_K; V_{2N})$ . First we suppose that  $a \equiv b \equiv 1 \pmod{2}$ . The equality of matrices:

$$(I - X)(I + X + \cdots + X^{p-1}) = 2I.$$

implies that the matrices  $I - X$  and  $I + X + \cdots + X^{p-1}$  are non-singular, i.e.,  $\text{rank}(I - X) = \text{rank}(I + X + \cdots + X^{p-1}) = 2N$ . By counting dimensions, we can see that the complex (4) is exact.

Next we consider the case that  $a \equiv b \equiv 0 \pmod{2}$ . In this case, all eigenvalues of  $X$  and  $Y$  are roots of unity. We divide this case into two parts:

- (i) either  $\text{SL}_{2N}(\mathbb{C})$ -matrices  $X$  or  $Y$  does not have the eigenvalue 1;  
(ii) both of  $X$  and  $Y$  have the eigenvalue 1.

In the case (i), we can assume that  $X$  does not have the eigenvalue 1. Since  $1 + \zeta + \cdots + \zeta^{p-1} = 0$  for any  $p$ -th root of unity  $\zeta \neq 1$ , we have that  $I + X + \cdots + X^{p-1} = O$ . Similarly we have that  $\text{rank}(I + Y + \cdots + Y^{q-1}) < 2N$ . Hence the rank of  $\partial_2$  is less than  $2N$ , which implies that  $H_2(W; V_{2N}) \neq \mathbf{0}$ .

Last we consider the case (ii). Without loss of generality, we can assume that

- the dimension of  $V_{2N}$  is greater than 2, i.e.,  $N \geq 2$ ;
- both of  $p$  and  $q$  are odd and;
- the pair  $p$  and  $a$  are coprime and;
- the pair  $q$  and  $b$  are also coprime.

The multiplicity  $\xi$  of eigenvalue 1 for  $X$  is given by twice the number of integers  $l$  such that  $p \leq p(2l-1) \leq 2N-1$ . Similarly we write  $\eta$  for the multiplicity of eigenvalue 1 for  $Y$ , which is given by twice the number of integers  $m$  such that  $q \leq q(2m-1) \leq 2N-1$ . By the explicit form of  $\partial_2$  as in (5), the kernel of  $\partial_2$  agrees with the intersection of  $\ker(I + X + \cdots + X^{p-1})$  and  $\ker(I + Y + \cdots + Y^{q-1})$ . Since the ranks of  $I + X + \cdots + X^{p-1}$  and  $I + Y + \cdots + Y^{q-1}$  are equal to  $\xi$  and  $\eta$ , we have that

$$\begin{aligned} \dim_{\mathbb{C}} \ker \partial_2 &= \dim_{\mathbb{C}} \ker(I + X + \cdots + X^{p-1}) + \dim_{\mathbb{C}} \ker(I + Y + \cdots + Y^{q-1}) \\ &\quad - \dim_{\mathbb{C}} (\ker(I + X + \cdots + X^{p-1}) \cup \ker(I + Y + \cdots + Y^{q-1})) \\ &\geq \dim_{\mathbb{C}} \ker(I + X + \cdots + X^{p-1}) + \dim_{\mathbb{C}} \ker(I + Y + \cdots + Y^{q-1}) - 2N \\ &= 2N - \xi - \eta \\ &\geq 2N - \frac{2N-1}{p} - 1 - \frac{2N-1}{q} - 1. \end{aligned}$$

We proceed to find the range on  $N$  satisfying that  $(2N-1)(1/p + 1/q) + 2 < 2N$ . We can rewrite the condition  $(2N-1)(1/p + 1/q) + 2 < 2N$  as

$$(6) \quad \frac{1}{1 - 1/p - 1/q} + 1 < 2N.$$

Since we assume that  $p \geq 3$  and  $q \geq 5$ , the inequality (6) holds for  $N \geq 2$ . Hence we have shown that  $\dim_{\mathbb{C}} \ker \partial_2 > 0$  under our assumptions, which implies that  $H_2(W; V_{2N})$  is nontrivial.  $\square$

#### 4. COMPUTATION OF HIGHER DIMENSIONAL REIDEMEISTER TORSION FOR TORUS KNOT EXTERIORS

Reidemeister torsion has the invariance under simple homotopy equivalences. In fact, a torus knot exterior  $E_K$  is simple homotopy equivalent to the 2-dimensional CW-complex

$W$  constructed as in Section 3. The Reidemeister torsion of  $E_K$  agrees with that of  $W$ . Therefore we can deduce the explicit value of Reidemeister torsion for a knot exterior  $E_K$  from the twisted chain complex (4) of  $W$ . We assume that  $\rho_{2N}$  defines an acyclic twisted chain complex  $C_*(E_K; V_{2N})$ . As we have seen in Section 3, such representation  $\rho_{2N}$  is given by an irreducible  $\mathrm{SL}_2(\mathbb{C})$ -representation of  $\pi_1(E_K)$  which satisfies that  $\rho(x^p) = \rho(y^q) = -I$ . We first show the explicit value of Reidemeister torsion for such  $\rho_{2N}$  and then consider the asymptotic behavior.

**4.1. Higher dimensional Reidemeister torsion for  $\rho_{2N}$ .** We write  $\{v_i \mid i = 1, \dots, 2N\}$  for a basis of  $V_{2N}$ . For the acyclic chain complex  $C_*(W; V_{2N})$ , we can set  $\tilde{\mathbf{b}}^i$  ( $i = 0, 1, 2$ ) as

$$\begin{aligned}\tilde{\mathbf{b}}^2 &= \{v_1 \otimes \tilde{e}^2, \dots, v_{2N} \otimes \tilde{e}^2\}, \\ \tilde{\mathbf{b}}^1 &= \{v_1 \otimes \tilde{e}_2^1, \dots, v_{2N} \otimes \tilde{e}_2^1\}\end{aligned}$$

and the others  $\tilde{\mathbf{b}}^i$  as  $\emptyset$ . The torsion of  $C_*(W; V_{2N})$  is given by the alternating product:

$$\begin{aligned}(7) \quad \prod_{i=0}^2 [\partial_{i+1} \tilde{\mathbf{b}}^{i+1} \cup \tilde{\mathbf{b}}^i / \mathbf{c}^i]^{(-1)^{i+1}} &= [\partial_3 \tilde{\mathbf{b}}^3 \cup \tilde{\mathbf{b}}^2 / \mathbf{c}^2]^{(-1)} \cdot [\partial_2 \tilde{\mathbf{b}}^2 \cup \tilde{\mathbf{b}}^1 / \mathbf{c}^1] \cdot [\partial_1 \tilde{\mathbf{b}}^1 \cup \tilde{\mathbf{b}}^0 / \mathbf{c}^0]^{(-1)} \\ &= \frac{\det(I + X + \dots + X^{p-1})}{\det(Y - I)} \\ &= \frac{2^{2N}}{\det(I - X) \det(Y - I)}.\end{aligned}$$

Here we use  $(I - X)(I + X + \dots + X^{p-1}) = I - X^p = 2I$ . The determinant  $\det(I - X)$  in the denominator of Eq. (7) is expressed as

$$\begin{aligned}(8) \quad \det(I - X) &= \prod_{k=1}^N (1 - e^{\frac{(2k-1)a\pi\sqrt{-1}}{p}})(1 - e^{-\frac{(2k-1)a\pi\sqrt{-1}}{p}}) \\ &= \prod_{k=1}^N 4 \sin^2\left(\frac{(2k-1)a\pi}{2p}\right)\end{aligned}$$

and  $\det(Y - I)$  is also expressed as

$$(9) \quad \det(Y - I) = \prod_{k=1}^N 4 \sin^2\left(\frac{(2k-1)b\pi}{2q}\right).$$

Substituting Eqs. (8) & (9) into (7), we obtain the explicit form of  $\mathrm{Tor}(C_*(W; V_{2N}))$  which agrees with the Reidemeister torsion of  $E_K$ .

**Proposition 4.1.** *If we choose an irreducible  $\mathrm{SL}_2(\mathbb{C})$ -representation  $\rho$  whose conjugacy class lies in  $\hat{R}_{a,b}(E_K)$  such that  $a \equiv b \equiv 1 \pmod{2}$ , then the Reidemeister torsion for  $E_K$  and  $\rho_{2N}$  is expressed as*

$$\mathrm{Tor}(E_K; \rho_{2N}) = \frac{2^{2N}}{\prod_{k=1}^N 4^2 \sin^2\left(\frac{(2k-1)a\pi}{2p}\right) \sin^2\left(\frac{(2k-1)b\pi}{2q}\right)}.$$

**4.2. Asymptotic behavior of Reidemeister torsions.** We consider the asymptotic behavior of  $\mathrm{Tor}(E_K; \rho_{2N})$  in increasing the dimension of representation  $\rho_{2N}$  to infinity. Menal-Ferrer and Porti [MFPa] showed the asymptotic behavior of the absolute value of Reidemeister torsion for hyperbolic knot exteriors  $M$ , which is expressed as

$$(10) \quad \lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M; \rho_{2N})|}{(2N)^2} = -\frac{\mathrm{Vol}(M)}{4\pi}.$$

Every torus knot exterior admits no hyperbolic structure and their hyperbolic volumes are zero. Our main theorem shows that the equation in (10) also holds for torus knot exteriors.

**Theorem 4.2.** *Under the assumption of Proposition 4.1, we have the following limit for the absolute value of Reidemeister torsion  $\text{Tor}(E_K; \rho_{2N})$ :*

$$(11) \quad \lim_{N \rightarrow \infty} \frac{\log |\text{Tor}(E_K; \rho_{2N})|}{(2N)^2} = 0.$$

*Proof of Theorem 4.2.* This follows from Proposition 4.1 and the existences on upper and lower bounds of  $|\sin((2k-1)a\pi/2p)|$  and  $|\sin((2k-1)b\pi/2q)|$  for  $1 \leq k \leq N$ , which are given dependently only on  $p$  and  $q$ . Since  $|\sin(\pi/2p)| \leq |\sin((2k-1)a\pi/2p)| \leq 1$  and  $|\sin(\pi/2q)| \leq |\sin((2k-1)b\pi/2q)| \leq 1$ , we have the following lower and upper bounds for  $|\text{Tor}(E_K; \rho_{2N})|$ :

$$\frac{2^{2N}}{4^{2N}} \leq |\text{Tor}(E_K; \rho_{2N})| \leq \frac{2^{2N}}{4^{2N} \sin^{2N}\left(\frac{\pi}{2p}\right) \sin^{2N}\left(\frac{\pi}{2q}\right)}.$$

We can see the asymptotic behavior of  $|\text{Tor}(E_K; \rho_{2N})|$  from the following inequalities:

$$\frac{\log |\text{Tor}(E_K; \rho_{2N})|}{(2N)^2} \geq \frac{-\log 2}{2N} \xrightarrow{N \rightarrow \infty} 0$$

and

$$\frac{\log |\text{Tor}(E_K; \rho_{2N})|}{(2N)^2} \leq \frac{-\log 2}{2N} - \frac{\log \left| \sin\left(\frac{\pi}{2p}\right) \sin\left(\frac{\pi}{2q}\right) \right|}{2N} \xrightarrow{N \rightarrow \infty} 0.$$

□

#### ACKNOWLEDGMENT

The author wishes to express his thanks to Joan Porti and Teruaki Kitano for drawing the author's attention to asymptotic behavior of higher dimensional Reidemeister torsion for torus knot exteriors. This work was motivated in RIMS Seminar "Representation spaces, twisted topological invariants and geometric structures of 3-manifolds". This research was supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

#### REFERENCES

- [Joh] D. Johnson, *A geometric form of Casson's invariant and its connection to Reidemeister torsion*, unpublished lecture notes.
- [KM] T. Kitano and T. Morifuji, *Twisted Alexander polynomials for  $sl(2; \mathbb{C})$ -irreducible representations of torus knots*, to appear in *Annali della Scuola Normale Superiore*.
- [MFPa] P. Menal-Ferrer and J. Porti, *Higher dimensional Reidemeister torsion invariants for cusped hyperbolic 3-manifolds*, arXiv:1110.3718.
- [MFPb] ———, *Twisted cohomology for hyperbolic three manifolds*, arXiv:1001.2242, to appear in *Osaka J. Math.*
- [Mül] W. Müller, *The asymptotics of the Ray–Singer analytic torsion of hyperbolic 3-manifolds*, arXiv:1003.5168v1.
- [Rag65] M. S. Raghunathan, *On the first cohomology of discrete subgroups of semisimple lie groups*, *Amer. J. Math.* **87** (1965), 103–139.
- [Tur01] V. Turaev, *Introduction to combinatorial torsions*, *Lectures in Mathematics*, Birkhäuser, 2001.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY 2-12-1 OOKAYAMA, MEGURO-KU TOKYO, 152-8551, JAPAN

*E-mail address:* shouji@math.titech.ac.jp